



# Heavy Metal Toxicity in Armed Conflicts Potentiates AMR in *A. baumannii* by Selecting for Antibiotic and Heavy Metal Co-resistance Mechanisms

## OPEN ACCESS

### Edited by:

Daniela Ceccarelli,  
European Commission, Belgium

### Reviewed by:

Zhi Ruan,  
Zhejiang University, China  
Max Mergeay,  
Belgian Nuclear Research Centre,  
Belgium

### \*Correspondence:

Antoine G. Abou Fayad  
aa328@aub.edu.lb  
Ghassan M. Matar  
gmatar@aub.edu.lb

†These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Antimicrobials, Resistance  
and Chemotherapy,  
a section of the journal  
Frontiers in Microbiology

Received: 09 July 2019

Accepted: 14 January 2020

Published: 03 February 2020

### Citation:

Bazzi W, Abou Fayad AG,  
Nasser A, Haraoui L-P, Dewachi O,  
Abou-Sitta G, Nguyen V-K, Abara A,  
Karah N, Landecker H, Knapp C,  
McEvoy MM, Zaman MH, Higgins PG  
and Matar GM (2020) Heavy Metal  
Toxicity in Armed Conflicts Potentiates  
AMR in *A. baumannii* by Selecting for  
Antibiotic and Heavy Metal  
Co-resistance Mechanisms.  
Front. Microbiol. 11:68.  
doi: 10.3389/fmicb.2020.00068

Wael Bazzi<sup>1,2,3†</sup>, Antoine G. Abou Fayad<sup>1,2,3\*†</sup>, Aya Nasser<sup>1,2,3</sup>, Louis-Patrick Haraoui<sup>4</sup>, Omar Dewachi<sup>5</sup>, Ghassan Abou-Sitta<sup>6</sup>, Vinh-Kim Nguyen<sup>7</sup>, Aula Abara<sup>8</sup>, Nabil Karah<sup>9</sup>, Hannah Landecker<sup>10</sup>, Charles Knapp<sup>11</sup>, Megan M. McEvoy<sup>12</sup>, Muhammad H. Zaman<sup>13</sup>, Paul G. Higgins<sup>14,15</sup> and Ghassan M. Matar<sup>1,2,3\*</sup>

<sup>1</sup> Department of Experimental Pathology, Immunology and Microbiology, Faculty of Medicine, American University of Beirut, Beirut, Lebanon, <sup>2</sup> Center for Infectious Diseases Research, American University of Beirut, Beirut, Lebanon, <sup>3</sup> World Health Organisation (WHO) Collaborating Center for Reference and Research on Bacterial Pathogens, Beirut, Lebanon, <sup>4</sup> Department of Microbiology and Infectious Diseases, University of Sherbrooke, Sherbrooke, QC, Canada, <sup>5</sup> Rutgers, The State University of New Jersey, Newark, NJ, United States, <sup>6</sup> Faculty of Medicine, American University of Beirut, Beirut, Lebanon, <sup>7</sup> The Graduate Institute of International and Developmental Studies, Geneva, Switzerland, <sup>8</sup> Department of Infection, Imperial College London, London, United Kingdom, <sup>9</sup> Department of Molecular Biology, Umea University, Umea, Sweden, <sup>10</sup> Department of Sociology and Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA, United States, <sup>11</sup> Civil and Environmental Engineering, University of Strathclyde, Glasgow, United Kingdom, <sup>12</sup> Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA, United States, <sup>13</sup> Department of Biomedical Engineering, Boston University, Boston, MA, United States, <sup>14</sup> Institute for Medical Microbiology, Immunology and Hygiene, University of Cologne, Cologne, Germany, <sup>15</sup> German Centre for Infection Research (DZIF), Partner Site Bonn-Cologne, Cologne, Germany

*Acinetobacter baumannii* has become increasingly resistant to leading antimicrobial agents since the 1970s. Increased resistance appears linked to armed conflicts, notably since widespread media stories amplified clinical reports in the wake of the American invasion of Iraq in 2003. Antimicrobial resistance is usually assumed to arise through selection pressure exerted by antimicrobial treatment, particularly where treatment is inadequate, as in the case of low dosing, substandard antimicrobial agents, or shortened treatment course. Recently attention has focused on an emerging pathogen, multi-drug resistant *A. baumannii* (MDRAb). MDRAb gained media attention after being identified in American soldiers returning from Iraq and treated in US military facilities, where it was termed “Iraqibacter.” However, MDRAb is strongly associated in the literature with war injuries that are heavily contaminated by both environmental debris and shrapnel from weapons. Both may harbor substantial amounts of toxic heavy metals. Interestingly, heavy metals are known to also select for antimicrobial resistance. In this review we highlight the potential causes of antimicrobial resistance by heavy metals, with a focus on its emergence in *A. baumannii* in war zones.

**Keywords:** *Acinetobacter baumannii*, bacteria, heavy metals, heavy metal tolerance, antimicrobial resistance, conflict, weapons

## INTRODUCTION

### Overview on Wars and Antimicrobial Resistance (AMR)

Over the past decades, *Acinetobacter baumannii* has emerged as a major driver of hospital-acquired Multi-Drug Resistant (MDR) infections (Manchanda et al., 2010; Williams et al., 2016; Mancilla-Rojano et al., 2019). In the decades following the Second World War, *A. baumannii* became one of the most prevalent pathogens during wars in Lebanon, Afghanistan, and Iraq, causing multiple outbreaks of MDR infections among combat casualties (Tong, 1972; CDC, 2004; Scott et al., 2007; Dallo and Weitao, 2010). According to The Centers for Disease Control and Prevention (CDC), *A. baumannii* was the single most isolated Gram-negative bacterium from war wounds during the recent wars in Iraq and Afghanistan and the number one causative agent of bloodstream infections among the US soldiers (CDC, 2004; Fournier et al., 2006). Moreover, the emergence of MDR, Extensively-Drug Resistant (XDR), and Pan-Drug Resistant (PDR) *A. baumannii* coincided with specific worldwide tension points such as the Lebanese Civil War (1975–1990) (Matar et al., 1992), Iraq-Iran war (1980–1988), Afghanistan war (2001–2014), Iraq war (2003–2011), and recently the Syrian war (2011–Present) (Tong, 1972; CDC, 2004; Scott et al., 2007). Warfare is associated with significant heavy metal contamination of the environment, due to destruction of built infrastructure and consequent release of HM and direct contamination from exploded ordnance and leakage from unexploded ordnance. HM resistance in bacteria is associated with antimicrobial resistance. In this review we investigate how heavy metal resistance can lead to antimicrobial resistance with a view to illuminating the emergence *A. baumannii*'s increased resistance in war regions.

### Living Systems and Heavy Metals

Heavy metals are a group of non-biodegradable metals and semi-metals (Metalloids) with high atomic weight and a density greater than 5 g/cm<sup>3</sup>. They have high electrical conductivity, malleability, metallic luster, and are capable of transferring electrons to form Cations (Jarup, 2003; Appenroth, 2010; Tchounwou et al., 2012). The major list of heavy metals includes: Cobalt (Co), Copper (Cu), Chromium (Cr), Zinc (Zn), Lead (Pb), Mercury (Hg), Arsenic (As), Cadmium (Cd), Nickel (Ni), Antimony (Sb), Boron (B), Barium (Ba), Silver (Ag), and Tungsten (W) (ATSDR, 2011). Essential heavy metals are required in many biological processes at low concentrations, where they serve as cofactors for metalloenzymes. For example, Cu serves as an integral cofactor for cytochrome c oxidase and superoxide dismutase required in mitochondrial electron transport and oxidative stress (Osredkar and Šuštar, 2011). Zn plays role in DNA and RNA polymerases catalytic functions (Markov et al., 1999). Ni and Cr are essential for the activity of urease and cytochrome P450 enzymes (Seiler and Berendonk, 2012). On the other hand, non-essential heavy metals such as Hg, Pb, Cd, and As have no known biological functions and are toxic even at low concentrations (ATSDR, 2011; Singh R. et al., 2011; Tchounwou et al., 2012).

Sources of heavy metals include water, soil, and rocks. Furthermore, they are present in the Earth's crust and date back to 4.5 billion years ago (Fru et al., 2016). The ecosystem is continuously flooded with high amounts of heavy metals due to volcanic eruptions, soil and rock erosion, industrial operations such as petroleum combustion and coal burning, in addition to agricultural activities that include pesticides and fungicides preparation (Seiler and Berendonk, 2012; Tchounwou et al., 2012; Jaroslawska and Piotrowska-Seget, 2014; Pal et al., 2017). Furthermore, heavy metals have been used in medical treatments as antimicrobial and anti-parasitic agents to treat skin infections and leishmaniasis respectively; anti-inflammatory compounds to treat itchiness, and in chemotherapy to treat cancer patients (Rizzotto, 2012; Ndagi et al., 2017; Pal et al., 2017).

### Heavy Metals-Mediated Bacterial Toxicity

Essential and non-essential heavy metals become toxic when exceeding specific concentrations that vary between different metals (Appenroth, 2010). In bacteria, toxic levels of heavy metals can accumulate in cells, altering cellular processes, inducing structural modifications, and ultimately leading to heavy metals-mediated damage (Rouch et al., 1995). Mechanisms of heavy metals toxicity include the production of Reactive Oxygen Species (ROS) that destroy essential biomolecules and sub-cellular organelles. When in excess, Pb, Cd, Cu, As, Ag, and Zn can induce oxidative damage in bacterial cells which leads to the production of free radicals that cause DNA damage and destabilize membranous integrity through lipid peroxidation (Szivák et al., 2009; Jaroslawska and Piotrowska-Seget, 2014; Williams et al., 2016). In addition, heavy metal ions can form complexes with thiol (R-SH)-containing enzymes to alter their functions. For example, Hg<sup>2+</sup>, Ag<sup>1+</sup>, and Cd<sup>2+</sup> can form covalent bonds with sulfhydryl functional groups (R-SH) present in enzymatic active sites thereby inducing structural conformational changes, and thus blocking their function (Rouch et al., 1995). Furthermore, heavy metals can act as competitive inhibitors leading to the displacement of essential ions from their target sites (Ianeva, 2009; Jan et al., 2015).

### Heavy Metals Resistance in Bacteria

Studies have shown that the presence of heavy metal resistance determinants is ubiquitous in almost all bacterial species (Silver and Phung le, 2005). For example, Farias et al. (2015), identified 35 bacterial strains from 8 different species harboring multi-metal resistant phenotypes from deep-sea hydrothermal vents. Moreover, *A. baumannii* isolated from agricultural soil and sediments, fuel-contaminated soil, and sewage water, have shown to exhibit resistance to various metals such as Hg, Ag, and As (Dhakephalkar and Chopade, 1994; Lima de Silva et al., 2012; Farias et al., 2015; Kim et al., 2015; El-Sayed, 2016; Huang et al., 2017). To date, six proposed mechanisms of heavy metal resistance have been elucidated: (1) Release of metal ions by extracellular barriers such as capsule, cell wall, and plasma membrane. (2) Extrusion of metal ions via efflux pumps or by diffusion. (3) Intracellular sequestration of metal

ions. (4) Extracellular sequestration of metal ions. (5) Bio-transformation/detoxification of toxic metal ions. (6) Decreased sensitivity of cellular targets to metal ions. In general, heavy metal resistance-encoding genes are carried on mobile genetic elements such as plasmids and transposons or on chromosomal DNA (Rouch et al., 1995; Silver and Phung le, 2005; Ianeva, 2009; Seiler and Berendonk, 2012; Hobman and Crossman, 2014). In the following sections, we will briefly discuss these mechanisms.

### Extrusion of Metal Ions via Efflux Pumps or Diffusion

Efflux pumps are the most prevalent bacterial tools conferring heavy metals resistance. This is achieved via Adenosine triphosphate (ATP) hydrolysis and/or through an electrochemical gradient of protons (Ianeva, 2009). Five major efflux system families are present in microorganisms: (1) ATP Binding Cassettes (ABC) family. (2) Resistance, Nodulation, Cell Division (RND) family. (3) Small Multi-Drug Resistance (SMR) family. (4) Multi-Drug and Toxic Compounds Efflux (MATE) transporters. (5) Major Facilitator Superfamily (MFS) family. These pumps differ in their amino acid sequence, substrate specificity, and energy consumption in pumping metal ions. For example, ATP Binding Cassettes (ABC) play a role in the efflux of metal ions and antimicrobial agents driven by ATP hydrolysis, while RNDs and SMRs pump out metal cations and antimicrobial agents via Chemiosmosis/Proton motive force (Silver and Phung le, 2005; Kourtesi et al., 2013; Abbas et al., 2017). Basal levels of efflux are usually not sufficient to confer heavy metal resistance in most bacterial species. However, changes in expression, either through mutations in promoter regions or efflux pump regulators, or inactivation of repressors, can lead to the over-expression of an efflux pump or confer resistance (Li et al., 2015; Blanco et al., 2016).

### Intracellular Sequestration of Metal Ions

One important resistance mechanism involves intracellular metal ions sequestration upon binding to metal ions binding proteins [Metallothioneins (MTs), Glutathione (GSH), and Metallochaperones] (Ianeva, 2009). For example, Ni can complex with  $\text{PO}_4^{3-}$ , leading to its intracellular precipitation. *Staphylococcus aureus*, *Providencia* spp., *Vibrio harveyi*, *Shewanella* spp., and *Bacillus* spp., can precipitate Pb as phosphate salts (Levinson et al., 1996; Smeaton et al., 2009; Shin et al., 2012). Cd, Hg, Ag, Pb, and Zn can be trapped on cysteine-rich MT polypeptides that provide tolerance to high concentrations of heavy metals. For example, *Synechococcus* spp., *Pseudomonas* spp., and *Anabaena* spp. tolerate high concentrations of heavy metals via MT trapping mechanisms (Olafson et al., 1988; Naik et al., 2012). Some bacteria use GSH as an alternative chelator to sequester metal ions. GSH scavenges and detoxifies metals via its thiol (R-SH) group. Lima et al. (2005), demonstrated the role of GSH in mediating tolerance to Cd in *Rhizobium leguminosarum*. Finally, metallochaperones like Cu chaperones (Cu<sup>1+</sup> binding chaperone CusF, Cu<sup>1+</sup>, and Cu<sup>2+</sup> periplasmic chaperones PcoC and PcoE) can bind, trap, and transport metal ions to metalloenzymes and thus, decrease their toxic effects and protect cellular compartments (Yang et al., 2010; Pal et al., 2017).

### Extracellular Sequestration of Metal Ions

In addition to intracellular sequestration, extracellular sequestration of heavy metals is an additional mechanism conferring bacterial resistance. This strategy provides a “Pre-defense” strategy as it occurs outside the bacterial cell. It involves the secretion of extracellular chelating proteins such as siderophores, oxalateoxalate, phosphate, and sulfide. However, this mechanism is mainly active in static environments when constant concentrations of heavy metals are present (Cunningham et al., 1993; Rouch et al., 1995). For example, *Streptomyces acidiscabies* can sequester Ni via hydroxamate siderophores while *Clostridium thermoaceticum* use sulfide to sequester Cd (Cunningham et al., 1993; Dimkpa et al., 2008).

### Bio-Transformation/Detoxification of Toxic Metal Ions

Enzymatic detoxification reduces metal toxicity, which is accomplished via oxidation, reduction, and methylation reactions (Rouch et al., 1995; Silver and Phung le, 2005). For example, Hg<sup>2+</sup> is reduced to a less toxic Hg<sup>0</sup> form via Mercury(II) reductase encoded by *merA* gene (Seiler and Berendonk, 2012; Pal et al., 2017). In addition, upon bacterial entry Cr<sup>6+</sup> is reduced to Cr<sup>3+</sup> while As<sup>3+</sup> is oxidized to As<sup>5+</sup> thus, decreasing their toxicity (Silver and Phung le, 2005). Interestingly, *Pseudomonas* spp., and *Acinetobacter* spp., induce Pb methylation to minimize its toxic effects. To date, Hg methylation has only been only documented in anaerobic bacteria (Parks et al., 2013; Jaroslawiecka and Piotrowska-Seget, 2014).

### Decreased Sensitivity of Cellular Targets to Metal Ions

Reducing sensitivity of cellular targets to metal ions is a way to minimize heavy metals toxicity. This is fulfilled via several mechanisms: (1) Decreasing bacterial susceptibility to metals by introducing mutations in resistance genes or determinants, or increasing the expression of the metal target site. (2) Producing a more resistant form of the metal target site upon activating an alternative target encoded on a plasmid. (3) Repairing DNA damage upon the activation of an SOS response which is the case of Cr-induced DNA lesions (Frohlich, 2013; Maret, 2015).

### Heavy Metals in Weapons

Explosives harbor huge amounts of Pb and Hg [Mercury(II) fulminate] (Navy U. S., 2008; Gebka et al., 2016). Zn, Cu, Ni, Pb, and Cr are used to coat bullets, missiles, gun barrels, and military vehicles (Audino, 2006; Casey, 2009). Ba, Sb, and B are weapon priming compounds, (Fitchett, 2019) and W is a kinetic bombardment due to its high density (19.3 g/cm<sup>3</sup>) (Rowlatt, 2014). In general, the use of heavy metals in weapons has increased since the end of World War II (Gebka et al., 2016).

### Known Resistance Mechanisms to Heavy Metals Frequently Found in Ordnance

#### Copper (Cu)

Copper (Cu) exists in nature as a free metallic element or alternates between 2 oxidative states Cu<sup>1+</sup> and Cu<sup>2+</sup>. In



humans, Cu is essential for blood vessel elasticity, brain development, maintenance of immune responses, and neurotransmitter production (Dorsey et al., 2004; Copper Development Association, 2018). Excess concentrations of Cu lead to kidney and liver damage, neurological, and immune diseases (Dorsey et al., 2004). In wars, Cu, Ni, Pb, and Cr are heavily used as coatings for bullets, missiles, gun barrels, and in military vehicles (tanks, trucks, and aircrafts). This could increase exposure to Cu in wartime and might explain increased observation of *A. baumannii* in studies conducted in these settings (Davis et al., 2005; Calhoun et al., 2008; Williams et al., 2016). Extensive studies on *Escherichia coli* and *Pseudomonas* spp. reveal four Cu homeostatic resistance systems: Cue, Cus, Pco, and Cop. Cue and Cus are chromosomally encoded efflux systems while Pco and Cop are plasmid encoded resistance systems (Pal et al., 2017).

### Cue System

The Cue system (Copper Efflux) is active at low Cu concentrations and under aerobic conditions. It consists of an inner membranous Cu<sup>1+</sup> exporting P-type ATPase (CopA) and a periplasmic multi-Cu Oxidase (CueO). *copA* and *cueO* are activated by a cytoplasmic transcriptional Cu-responsive Regulon (CueR) upon sensing increased Cu concentrations (Hobman and Crossman, 2014; Delmar et al., 2015; Pal et al., 2017; **Figure 1**).

### Cus System

Unlike the Cue system, the Cus efflux system is active at high Cu concentrations, is strictly anaerobic, and pumps out Cu and Ag cations (Silver and Phung le, 2005; Singh S.K. et al., 2011; Delmar et al., 2015). It detoxifies Cu in the periplasmic compartment, unlike the Cue system that extrudes periplasmic and cytoplasmic Cu (Jaroslawska and Piotrowska-Seget, 2014). The Cus system consists of 4 genes forming the *cusCFBA* operon, which is regulated by a two-component regulatory system (CusS/CusR) (Franke et al., 2003; Silver and Phung le, 2005; Pal et al., 2017). CusS, a histidine kinase, is activated upon Cu/Ag stimulation, while CusR is a DNA-binding transcriptional activator that activates *cusCFBA* expression. CusC, CusB, and CusA form a multi-Cu/Ag efflux pump (CusCBA) that functions as a proton-antiporter. Cu<sup>1+</sup>/Ag<sup>1+</sup> are transported to CusCBA via the periplasmic metallo-chaperone, CusF (Franke et al., 2003; Silver and Phung le, 2005; Pal et al., 2017). Cu resistance via the Cue and Cus systems are detailed in **Figure 1**.

### Pco System

The *E. coli*-resistant Pco system (Plasmid-borne-Copper Resistance) found in Cu-fed pigs consists of two operons, *pcoGFE*, and *pcoABCDS* that are encoded by a 9-10 gene cluster (Brown et al., 1995). Like the Cus system, the Pco system is regulated by a two-component regulatory system (PcoR/PcoS) (Brown et al., 1995; Pal et al., 2017). To actively function, the Pco system requires the action of CopA from the Cue system in addition to PcoA and PcoC. PcoA is a multi-Cu Oxidase that oxidizes Cu<sup>1+</sup> to Cu<sup>2+</sup>, while PcoC is a periplasmic Cu-binding protein that acts as a chaperone which delivers

Cu<sup>1+</sup> to CopA during oxidation and to PcoD. PcoD is an inner membrane Cu transporter that is involved in Cu uptake. PcoB and PcoE are an outer membrane transporter and a metallo-chaperone, respectively (**Figure 2**; Lee et al., 2002; Bondarczuk and Piotrowska-Seget, 2013; Pal et al., 2017).

### Cop System

The Cop system is encoded by a cluster of 6 plasmid-borne genes arranged in two operons, *copABCD* and *copRS* (Pal et al., 2017). *copABCD* and *copRS* are homologs of *pcoABCDS*. The Cop determinants are genetically associated with the Cop system and have similar roles. *copABCD* is under the regulation of CopR/CopS. Protein products are associated with Cu sequestration in the periplasm and outer membrane (Bondarczuk and Piotrowska-Seget, 2013; Pal et al., 2017). In *Cupriavidus metallidurans* CH34 and *E. coli*, Cu ions can be sequestered by CusF in the periplasm, exported by the RND-driven CusCBA efflux pump, or oxidized to Cu<sup>2+</sup> (Nies, 2016).


### Mercury (Hg)

Mercury (Hg) is released into the environment via geological and human activities such as soil and rock erosion, volcanic eruptions, mining, and fuel combustion (Tchounwou et al., 2012). In the wake of recent conflicts in Lebanon, Syria, Iraq, Yemen, and Afghanistan, the Middle East has become one of the most polluted regions with Hg (Gworek et al., 2017). This may have led to an increased bacterial tolerance to this metal. In humans, Hg has no biological role and at very low concentrations it is fatal, leading to brain, lung, and kidney failure (Risher and Rob, 1999).

Hg can access bacteria in 2 forms: organic (CH<sub>3</sub>-Hg<sup>+</sup>) and inorganic (Hg<sup>2+</sup>), both of which are toxic (Hobman and Crossman, 2014). Despite this toxicity, several bacterial species have developed resistance mechanisms to CH<sub>3</sub>-Hg<sup>+</sup>/Hg<sup>2+</sup> via the *mer* operon, and are found mainly in war zone regions (Mirzaei et al., 2008; Pérez-Valdespino et al., 2013; Hobman and Crossman, 2014). The *mer* operon is present on plasmids and transposons and consists of a cluster of 8 genes *merTPCAGBDE* regulated by MerR (Silver and Phung le, 2005; Hobman and Crossman, 2014). This operon encodes a chain of proteins that bind CH<sub>3</sub>-Hg<sup>+</sup>/Hg<sup>2+</sup> and oxidize them, such as MerA [Mercury(II) Reductase], the key player in Hg<sup>2+</sup> detoxification (**Figure 3**).

### Arsenic (As)

Arsenic (As) is released into the environment from soil and rock erosion, volcanic eruptions, mining, and crops treated with pesticides and herbicides (Paez-Espino et al., 2009; Tchounwou et al., 2012). In 1918, two organic As compounds, Lewisite (C<sub>2</sub>H<sub>2</sub>AsCl<sub>3</sub>) and Adamsite (C<sub>12</sub>H<sub>9</sub>AsClN) were developed by the US army as chemical weapons; both are classified as potential bioterrorism agents by CDC (2013). Agent Blue, an arsenical mixture of cacodylic acid and sodium cacodylate was sprayed by the United States on crops as part of “resource deprivation” strategies in the Vietnam war beginning in 1962 (Radke et al., 2014). The use of chemical weapons in the Syrian Civil War has been confirmed by the United Nations. This resulted in increased bacterial resistance to As via oxidation, reduction,



methylation, efflux, and intracellular sequestration on cysteine-rich peptides (Silver and Phung le, 2005; Paez-Espino et al., 2009), and was associated with detrimental health effects ranging from cardiovascular disease, respiratory disorders, gastro-intestinal symptoms, hematological disorders, diabetes, neurological, and developmental anomalies (Chou et al., 2007; Tchounwou et al., 2012).


As exists in two chemical forms: inorganic and organic. Inorganic As occurs as pentavalent Arsenate ( $\text{As}^{5+}$ ), trivalent Arsenite ( $\text{As}^{3+}$ ), elemental Arsenic ( $\text{As}^0$ ), and Arsenide ( $\text{As}^{3-}$ ) with  $\text{As}^{3+}$  and  $\text{As}^{5+}$  being the most toxic inorganic forms and most prevalent in nature (Nies, 1999; Paez-Espino et al., 2009; Tchounwou et al., 2012). Organic As is less toxic than inorganic arsenicals (Chou et al., 2007). Bacterial resistance to As is mainly encoded by efflux via the *ars* operon, which can be plasmid or chromosomally driven, even though it can also be encoded by other genetic determinants such as *arr* genes and *aox* genes (Figure 4; Silver and Phung le, 2005; Paez-Espino et al., 2009).

### Chromium (Cr)

Chromium (Cr) is the 7th most abundant heavy metal in the earth's crust and is present in nature in several oxidation states

ranging from divalent (+2) to hexavalent (+6)  $\text{Cr}^{3+}$  and  $\text{Cr}^{6+}$  are the most stable. While  $\text{Cr}^{3+}$  is naturally present in the environment,  $\text{Cr}^{6+}$  is mostly produced by industrial processes such as mining, electroplating, dye production, and leather tanning (Ahemad, 2014; Joutey et al., 2015; Pradhan et al., 2016). In weapons, Cr was initially used by the Chinese to coat metal weapons (Dunham, 2019). Nowadays, Cr is heavily used to coat gun barrels, where it is used as a bore protection (Audino, 2006). Moreover, Cr levels were highest in deciduous teeth from Iraqi patients during the Iraqi war, which highlights the heavily polluted Middle Eastern region with heavy metals (Savabieasfahani et al., 2016). The solubility and oxidizing potential of  $\text{Cr}^{6+}$  makes it 1000× more toxic to humans as compared to  $\text{Cr}^{3+}$ , and this makes it a strong factor associated with nasal and bronchogenic carcinomas (International Agency for Research on Cancer [IARC], 1990; Ahemad, 2014).

In bacteria, Cr has no metabolic role and thus, it is toxic in several species such as *Pantoea* spp., *Aeromonas* spp., *Acinetobacter* spp., and *E. coli* (Nies, 1999). However, many bacteria developed Cr resistance via 5 reported mechanisms that are mostly plasmid encoded (Camargo et al., 2005; Joutey et al., 2015). (1) Reduction of  $\text{Cr}^{6+}$  uptake. (2)  $\text{Cr}^{6+}$  efflux.



(3) Activation of oxidative stress related enzymes. (4) Repairing DNA damage induced by  $\text{Cr}^{6+}$  and its derivatives. (5)  $\text{Cr}^{6+}$  reduction (Figure 5; Branco et al., 2008; Panda and Sarkar, 2012; Ahemad, 2014; Joutey et al., 2015; Pradhan et al., 2016).

### Lead (Pb)

Lead (Pb) is predominantly released into the environment from human activities such as manufacturing pipes, X-ray shields, lead-acid storage batteries, munitions, and bullets (Abadin et al., 2007; Jaroslawska and Piotrowska-Seget, 2014). It exists in two main oxidative states ( $\text{Pb}^{2+}$  and  $\text{Pb}^{4+}$ ). In addition to bullets, Pb is present in explosives that ignite gunpowder. It usually vaporizes upon firing and thus, Pb fumes and dust are inhaled, leading to brain damage, anemia, and high blood pressure (Abadin et al., 2007; Dermatas and Chrysochoou, 2007).


Pb toxicity involves inducing cellular damage through ROS formation, disrupting enzymatic conformations, and interfering in calcium (Ca) metabolism (Tchounwou et al., 2012). Due

to the widespread Pb contamination, bacteria have developed Pb resistance mechanisms (Levinson et al., 1996; Jaroslawska and Piotrowska-Seget, 2014). (1) Adsorption of Pb on EPS and bacterial cell wall. (2) Reducing Pb accumulation via intracellular and extracellular precipitation. (3) Pb sequestration via intracellular proteins. (4) Pb detoxification via methylation. (5) Pb extrusion via efflux pumps (Figure 6; Levinson et al., 1996; Silver and Phung le, 2005; Jaroslawska and Piotrowska-Seget, 2014; Subramanian, 2018).

### Association Between Heavy Metals and AMR

Worldwide concerns about heavy metal contamination, resistance, and its ability to induce AMR are increasing. These concerns are associated with the heavy metals used in manufacturing weapons; most heavy metals are non-biodegradable and persist in the environment. Moreover, many






bacterial species evolved resistance mechanisms to combat metals toxicity (Seiler and Berendonk, 2012; Yu et al., 2017). These mechanisms are encoded by resistant genes to heavy metals and antimicrobial agents that are physically linked on mobile genetic elements (Seiler and Berendonk, 2012; Yu et al., 2017). More importantly, heavy metals can induce selective pressure on microbial populations leading to antimicrobial resistance through a mechanism called “co-selection” which occurs via 3 major ways (Seiler and Berendonk, 2012).

### Co-resistance

Co-resistance occurs when genes encoding resistance to heavy metals and antimicrobial agents are physically linked/located

in close proximity to each other on mobile genetic elements such as plasmids, genomic islands (GIs), transposons, or integrons (Trevors and Oddie, 1986; Seiler and Berendonk, 2012; Yu et al., 2017). For example, in Cu-resistant *Enterococcus faecium* isolated from pigs, *tcrB* and genes encoding resistance to erythromycin and vancomycin are encoded on the same conjugative plasmid (Hasman and Aarestrup, 2002; Silveira et al., 2014). Moreover, in *Serratia marcescens*, plasmid-borne resistance to chloramphenicol, kanamycin, and tetracycline is genetically linked to As, Cu, Hg, and Ag resistance genes (Gilmour et al., 2004). Interestingly, Whole-Genome Sequencing (WGS) analysis in *Salmonella typhi* reveal a genetic association between Hg resistance and several unrelated antimicrobial agents



**FIGURE 4 |** Bacterial resistance to arsenic. (As) enter the bacterial cell using Phosphate-Specific Transporters (Pst) and Type III Transporters (PIT) in the case of  $As^{5+}$  and Aqueglycerolporins in the case of  $As^{3+}$  (Paez-Espino et al., 2009). The *ars* operon harbors 3 co-transcribed core genes that confer resistance not only to  $As^{3+}$  and  $As^{5+}$ , but also to Antimony ( $Sb^{3+}$ ). *arsR* encodes a Transcriptional Repressor, *arsC* encodes a Cytoplasmic Arsenite Reductase, and *arsB* encodes a membrane bound Arsenite Efflux Pump. Two additional genes may be present within the *ars* operon, *arsA* and *arsD*. The former encodes an intracellular ATPase which binds *ArsB* to form an *ArsA-ArsB* ATPase Efflux Pump, while the latter is a metallochaperone that binds and delivers  $As^{3+}$  and  $Sb^{3+}$  to *ArsA-ArsB* complex for efflux, in addition to its role as a trans-activating co-repressor of the *ars* operon along with *ArsR* (Silver and Phung le, 2005; Paez-Espino et al., 2009; Hobman and Crossman, 2014). Moreover, some microorganisms escape As toxicity by methylation thus, leading to the production of less toxic and volatile derivatives that diffuse outside the bacterial cell (Paez-Espino et al., 2009). Besides As toxicity, bacteria belonging to the *Shewanella* spp., *Sulfurospirillum* spp., *Clostridium* spp., and *Bacillus* spp., use  $As^{5+}$  as a final electron acceptor during anaerobic respiration by reducing it to  $As^{3+}$ , while other bacteria use  $As^{3+}$  as an electron donor and oxidize it to  $As^{5+}$  during aerobic oxidation (Paez-Espino et al., 2009). The oxidation/reduction processes are mediated by the Respiratory Arsenate Reductase and Respiratory Arsenite Oxidase that are encoded by the *arrAB* operon and *asoAB* genes respectively (Silver and Phung le, 2005; Paez-Espino et al., 2009). Adapted and modified with permission from Paez-Espino et al. (2009).


resistance genes (chloramphenicol, ampicillin, streptomycin, sulfonamide, and trimethoprim) (Wireman et al., 1997).

## Cross-Resistance

Cross-resistance occurs when one resistance mechanism confers resistance to heavy metals and antimicrobial agents simultaneously. This is mainly achieved via multi-drug efflux pumps (Baker-Austin et al., 2006; Pal et al., 2017). The MdrL efflux pump in *Listeria monocytogenes* encodes resistance to Zn, Co, Cr, erythromycin, josamycin, and clindamycin (Mata et al., 2000). Moreover, the DsbA-DsbB (Disulfide Bond) multi-drug efflux system in *Burkholderia cepacia* induces cross-resistance to  $\beta$ -lactams, kanamycin, erythromycin, novobiocin, ofloxacin, and  $Zn^{2+}$  and  $Cd^{2+}$  metal ions (Hayashi et al., 2000). In addition, resistance to antimicrobial agents, Co, and Cu is mediated via the CmeABC multi-drug efflux pump in *Campylobacter jejuni* (Lin et al., 2002).

## Co-regulatory Resistance

Co-regulation is the least common mechanism of co-selection. It is fulfilled when resistant genes to antimicrobial agents and




**FIGURE 5 |** Bacterial resistance to chromium. Five Cr resistance mechanisms are reported (Camargo et al., 2005; Joutey et al., 2015). (1) Reduction of  $Cr^{6+}$  uptake.  $Cr^{6+}$  exists in the form of Oxyanions Chromate ( $CrO_4^{2-}$ ) and Dichromate ( $Cr_2O_7^{2-}$ ). Bacterial cells can reduce  $Cr^{6+}$  uptake via the sulfate transport system (Nies, 1999; Ahemad, 2014; Joutey et al., 2015). (2)  $Cr^{6+}$  efflux. Studies reveal that *P. aeruginosa* and *Alcaligenes eutrophus* can extrude  $Cr^{6+}$  by active efflux through *ChrA* (Chromate) pump (Collard et al., 1994; Alvarez et al., 1999). In 2008, a plasmid encoded operon (*chrBACF*) was identified in *Ochrobactrum tritici* responsible for Cr efflux, where *chrB* and *chrA* are the main genes involved (Branco et al., 2008). (3) Activation of oxidative stress related enzymes. When  $Cr^{6+}$  enter the bacterial cell, it interacts with reducing agents such as Nicotinamide Adenine Dinucleotide Phosphate (NADPH) and Ascorbic Acid to produce free radicals and unstable Cr intermediates ( $Cr^{4+}$  and  $Cr^{5+}$ ) that are further reduced to  $Cr^{3+}$ . End products of these reactions cause oxidative stress leading to protein and DNA damage. This induces the up-regulation of antioxidants enzymes that scavenge ROS and protect cellular compartments (Ahemad, 2014; Joutey et al., 2015; Pradhan et al., 2016). (4) Repairing DNA damage induced by  $Cr^{6+}$  and its derivatives. This is achieved via SOS response activation. Several studies highlight the roles of RuvAB, RecA, and RecG (helicases) in mediating Cr resistance through repairing  $Cr^{6+}$  induced DNA damage (Miranda et al., 2005; Morais et al., 2011). (5)  $Cr^{6+}$  reduction.  $Cr^{6+}$  can be reduced aerobically or anaerobically to a less toxic form  $Cr^{3+}$ . Aerobic reduction uses cytoplasmic soluble reductases and NADPH, while anaerobic reduction uses membrane reductases belonging to the electron transport chain (cytochromes b and c, and hydrogenases) (Morais et al., 2011; Ahemad, 2014; Joutey et al., 2015). Adapted and modified with permission from Ahemad (2014).

heavy metals are controlled by a mutual regulatory protein (Pal et al., 2017). A very well characterized co-regulatory resistance system is the CzcS-CzcR two component regulatory system of *P. aeruginosa*. This system induces resistance to  $Zn^{2+}$ ,  $Cd^{2+}$ , and  $Co^{2+}$  by activating the expression of *czcCBA* (Cobalt Zinc Cadmium) efflux pump and to the carbapenem imipenem by suppressing expression of the *OprD* porin encoding gene (Perron et al., 2004).

## Whole-Genome Sequencing and Heavy Metal Resistance

Many reports highlighted the importance of WGS as an effective tool to detect genome-wide modifications and the emergence of heavy metal resistance genes. Nowadays, it is feasible to assess the entire bacterial genome at low costs and in a timely manner, making it an ideal method for AMR surveillance. Therefore, WGS





provides a practical solution to evaluate genomes and determine resistant genes for compounds that are not frequently assessed. Moreover, this tool allows scientists to discover novel resistance mechanisms and provides valuable information to researchers and clinicians in antimicrobial prescriptions (Khromykh and Solomon, 2015; Jagadeesan et al., 2019). For example, the arsenic resistance cassette, *arsRCDAB*, present on a class 1 integron and mobilized on a conjugative plasmid was detected in two *Salmonella enterica* isolates from Singapore with high tolerance to arsenate (Wilson et al., 2019). Moreover, heavy metal resistant genes to Manganese ( $Mn^{2+}$ ) and  $Cd^{2+}$  in addition to exopolysaccharides production (EPS) were documented upon WGS analysis in *Pseudaminobacter manganicus* isolated from a manganese mine. This aspect sheds light on metal removal/adsorption and reflects bioremediation capabilities in

contaminated regions (Xia et al., 2018). In *A. baumannii*, WGS analysis revealed the role of two integrases in the excision and circularization of heavy metal resistance (GIs) in *E. coli* (Al-Jabri et al., 2018). Bacterial WGS and typing databases such as BacWGSTdb (PMID: 26433226)<sup>1</sup> serve as easy, rapid, powerful, and convenient tools to assess AMR and provide valuable information to WGS analysis and daily use in the clinical microbiology laboratories (Ruan and Feng, 2016). Moreover, “BacMet: Antibacterial Biocide and Metal Resistance Genes Database” provides an accurate and high quality resource of bacterial genes associated with heavy metal resistance present in literature<sup>2</sup>. These available databases serve as important tools to help understand bacterial resistance mechanisms to heavy metals by linking various factors and parameters involved.

## CONCLUSION

The reason for the rapid emergence of drug resistant *A. baumannii* in war-wounded patients remains unclear. Heavy metal contaminated areas may be driving the increase in antimicrobial resistance. This may explain an observed increase in bacterial resistance to both heavy metals and antimicrobial agents and lead to the development of novel mechanisms of resistance. The role of this pathway in *A. baumannii* is poorly understood.

Until now antimicrobial resistance has been largely attributed to poor antimicrobial stewardship in humans and in animals. The mechanisms described above, whereby heavy metals may produce antimicrobial resistance in their absence, identifies a potential pathway driving global antimicrobial resistance that would not be addressed through improved antimicrobial stewardship. This pathway would be facilitated in wartime, and could explain the emergence of previously little reported pathogens as possible amplification points along this pathway.

Further research into the role of heavy metals driving antimicrobial resistance, its influence in war zones, and the contribution of *A. baumannii* as a reservoir, are therefore warranted given the implications for addressing the global AMR crisis.

## AUTHOR CONTRIBUTIONS

AGA, WB, AN, L-PH, MZ, and PH contributed to reviewing the literature and the write up of the manuscript. V-KN, HL, OD, GA-S, MM, NK, AA, CK, and GM contributed to the editing of the manuscript.

## FUNDING

This project was part of the War, AntiMicrobial Resistance, and *Acinetobacter baumannii* (WAMRA) consortium and was supported by the Medical Practice Plan, Faculty of Medicine, American University of Beirut.

<sup>1</sup><http://bacdb.org/BacWGSTdb>

<sup>2</sup><http://bacmet.biomedicine.gu.se/>

## REFERENCES

- Abadin, H., Ashizawa, A., Stevens, Y.-W., ATSDR, Lladós, F., Diamond, G., et al. (2007). *Toxic Substances Portal- Lead*. Atlanta: Agency for Toxic Substances and Disease Registry. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24049859>
- Abbas, S. Z., Rafatullah, M., Hossain, K., Ismail, N., Tajarudin, H. A., and Abdul Khalil, H. P. S. (2017). A review on mechanism and future perspectives of cadmium-resistant bacteria. *Int. J. Environ. Sci. Technol.* 15, 243–262. doi: 10.1007/s13762-017-1400-5
- Ahemad, M. (2014). Bacterial mechanisms for Cr(VI) resistance and reduction: an overview and recent advances. *Folia Microbiol.* 59, 321–332. doi: 10.1007/s12223-014-0304-8
- al-Aoukaty, A., Appanna, V. D., and Huang, J. (1991). Exocellular and intracellular accumulation of lead in *Pseudomonas fluorescens* ATCC 13525 is mediated by the phosphate content of the growth medium. *FEMS Microbiol. Lett.* 67, 283–290. doi: 10.1111/j.1574-6968.1991.tb04478.x
- Al-Jabri, Z., Zamudio, R., Horvath-Papp, E., Ralph, J. D., Al-Muharrami, Z., Rajakumar, K., et al. (2018). Integrase-controlled excision of metal-resistance genomic islands in *Acinetobacter baumannii*. *Genes* 9:E366. doi: 10.3390/genes9070366
- Alvarez, A. H., Moreno-Sanchez, R., and Cervantes, C. (1999). Chromate efflux by means of the ChrA chromate resistance protein from *Pseudomonas aeruginosa*. *J. Bacteriol.* 181, 7398–7400. doi: 10.1128/jb.181.23.7398-7400.1999
- Appenroth, K.-J. (2010). What are “heavy metals” in plant sciences? *Acta Physiol. Plant.* 32, 615–619. doi: 10.1007/s11738-009-0455-4
- ATSDR (2011). *Toxic Substances Portal*. Available at: <https://www.atsdr.cdc.gov/> (accessed March 22, 2019).
- Audino, M. J. (2006). *Use of Electroplated Chromium in Gun Barrels*. Washington, DC: Army Benet Laboratories.
- Baker-Austin, C., Wright, M. S., Stepanauskas, R., and McArthur, J. V. (2006). Co-selection of antibiotic and metal resistance. *Trends Microbiol.* 14, 176–182. doi: 10.1016/j.tim.2006.02.006
- Blanco, P., Hernando-Amado, S., Reales-Calderon, J. A., Corona, F., Lira, F., Alcalde-Rico, M., et al. (2016). Bacterial multidrug efflux pumps: much more than antibiotic resistance determinants. *Microorganisms* 4:E14. doi: 10.3390/microorganisms4010014
- Bondarczuk, K., and Piotrowska-Seget, Z. (2013). Molecular basis of active copper resistance mechanisms in Gram-negative bacteria. *Cell Biol. Toxicol.* 29, 397–405. doi: 10.1007/s10565-013-9262-1
- Boyd, E. S., and Barkay, T. (2012). The mercury resistance operon: from an origin in a geothermal environment to an efficient detoxification machine. *Front. Microbiol.* 3:349. doi: 10.3389/fmicb.2012.00349
- Branco, R., Chung, A. P., Johnston, T., Gurel, V., Morais, P., and Zhitkovich, A. (2008). The chromate-inducible chrBACF operon from the transposable element TnOtChr confers resistance to chromium(VI) and superoxide. *J. Bacteriol.* 190, 6996–7003. doi: 10.1128/JB.00289-08
- Brown, N. L., Barrett, S. R., Camakaris, J., Lee, B. T., and Rouch, D. A. (1995). Molecular genetics and transport analysis of the copper-resistance determinant (pco) from *Escherichia coli* plasmid pRJ1004. *Mol. Microbiol.* 17, 1153–1166. doi: 10.1111/j.1365-2958.1995.mmi\_17061153.x
- Calhoun, J. H., Murray, C. K., and Manring, M. M. (2008). Multidrug-resistant organisms in military wounds from Iraq and Afghanistan. *Clin. Orthop. Relat. Res.* 466, 1356–1362. doi: 10.1007/s11999-008-0212-9
- Camargo, F. A. O., Okeke, B. C., Bento, F. M., and Frankenberger, W. T. (2005). Diversity of chromium-resistant bacteria isolated from soils contaminated with dichromate. *Appl. Soil Ecol.* 29, 193–202. doi: 10.1016/j.apsoil.2004.10.006
- Casey, T. (2009). *US Military Targets Toxic Enemy #1: Hexavalent Chromium*. Available at: <https://cleantechnica.com/2009/07/04/us-military-targets-toxic-enemy-1-hexavalent-chromium/> (accessed July 4, 2009).
- CDC (2004). *Acinetobacter baumannii* infections among patients at military medical facilities treating injured U.S. service members, 2002–2004. *MMWR Morb. Mortal. Wkly. Rep.* 53, 1063–1066.
- CDC (2013). *Centers for Disease Control and Prevention. The Emergency Response Safety and Health Database. Lewisite (L): Blister Agent*. Available at: [https://www.cdc.gov/niosh/ershdb/emergencyresponsecard\\_29750006.html](https://www.cdc.gov/niosh/ershdb/emergencyresponsecard_29750006.html) (accessed April 01, 2019).
- Chou, S., Harper, C., ATSDR, Ingeman, L., Lladós, F., Colman, J., et al. (2007). *Toxic Substances Portal- Arsenic*. Available at: <https://www.atsdr.cdc.gov/toxprofiles/tp2.pdf>
- Collard, J. M., Corbisier, P., Diels, L., Dong, Q., Jeanthou, C., Mergeay, M., et al. (1994). Plasmids for heavy metal resistance in *Alcaligenes eutrophus* CH34: mechanisms and applications. *FEMS Microbiol. Rev.* 14, 405–414.
- Copper Development Association (2018). *Benefits of Copper*. Available at: <https://copperalliance.org.uk/benefits-copper/health/> (accessed March 22, 2019).
- Cunningham, D. P., Leon, L., and Lundie, J. R. (1993). Precipitation of cadmium by *Clostridium thermoaceticum*. *Appl. Environ. Microbiol.* 59, 7–14. doi: 10.1128/aem.59.1.7-14.1993
- Dallo, S. F., and Weita, T. (2010). Insights into acinetobacter war-wound infections, biofilms, and control. *Adv. Skin Wound Care* 23, 169–174. doi: 10.1097/01.ASW.0000363527.08501.a3
- Davis, K. A., Moran, K. A., McAllister, C. K., and Gray, P. J. (2005). Multidrug-resistant acinetobacter extremity infections in soldiers. *Emerg. Infect. Dis.* 11, 1218–1224.
- Delmar, J. A., Su, C.-C., and Yu, E. W. (2015). Heavy metal transport by the CusCFBA efflux system. *Protein Sci.* 24, 1720–1736. doi: 10.1002/pro.2764
- Dermatas, D., and Chrysochoou, M. (2007). Lead particle size and its association with firing conditions and range maintenance: implications for treatment. *Environ. Geochem. Health* 29, 347–355. doi: 10.1007/s10653-007-9092-2
- Dhakephalkar, P. K., and Chopade, B. A. (1994). High levels of multiple metal resistance and its correlation to antibiotic resistance in environmental isolates of acinetobacter. *Biometals* 7, 67–74.
- Dimkpa, C., Svatos, A., Merten, D., Buchel, G., and Kothe, E. (2008). Hydroxamate siderophores produced by *Streptomyces acidiscabies* E13 bind nickel and promote growth in cowpea (*Vigna unguiculata* L.) under nickel stress. *Can. J. Microbiol.* 54, 163–172. doi: 10.1139/w07-130
- Dorsey, A., Lisa, I., and Steven, S. (2004). *Toxic Substances Portal - Copper*. Available at: <https://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=206&tid=37> (accessed September 2004).
- Dunham, W. (2019). *Scientists Solve Mystery of Pristine Weapons of China's Terracotta Warriors*. Available at: <https://www.reuters.com/article/us-science-warriors/scientists-solve-mystery-of-pristine-weapons-of-chinas-terracotta-warriors-idUSKCN1RG2HT> (accessed April 5, 2019).
- El-Sayed, M. H. (2016). Multiple heavy metal and antibiotic resistance of *Acinetobacter baumannii* strain HAF-C 13 isolated from industrial effluents. *Am. J. Microbiol. Res.* 4, 26–36.
- Farias, P., Espirito Santo, C., Branco, R., Francisco, R., Santos, S., Hansen, L., et al. (2015). Natural hot spots for gain of multiple resistances: arsenic and antibiotic resistances in heterotrophic, aerobic bacteria from marine hydrothermal vent fields. *Appl. Environ. Microbiol.* 81, 2534–2543. doi: 10.1128/AEM.03240-14
- Fitchett, B. (2019). *Priming Compounds and Primers Introduction*. *Bev Fitchett's guns magazine*. Available at: <https://www.bevfitchett.us/ballistics/priming-compounds-and-primers-introduction.html>
- Fournier, P. E., Richet, H., and Weinstein, R. A. (2006). The epidemiology and control of *Acinetobacter baumannii* in health care facilities. *Clin. Infect. Dis.* 42, 692–699. doi: 10.1086/500202
- Franke, S., Grass, G., Rensing, C., and Nies, D. H. (2003). Molecular analysis of the copper-transporting efflux system CusCFBA of *Escherichia coli*. *J. Bacteriol.* 185, 3804–3812. doi: 10.1128/jb.185.13.3804-3812.2003
- Frohlich, E. (2013). Cellular targets and mechanisms in the cytotoxic action of non-biodegradable engineered nanoparticles. *Curr. Drug Metab.* 14, 976–988. doi: 10.2174/1389200211314090004
- Fru, E. C., Nathalie, R., Camille, P., Stefan, L., Per, A., Dominik, W., et al. (2016). Cu isotopes in marine black shales record the great oxidation event. *Proc. Natl. Acad. Sci. U.S.A.* 113, 4941–4946. doi: 10.1073/pnas.1523544113
- Gebka, K., Beldowski, J., and Beldowska, M. (2016). The impact of military activities on the concentration of mercury in soils of military training grounds and marine sediments. *Environ. Sci. Pollut. Res. Int.* 23, 23103–23113. doi: 10.1007/s11356-016-7436-0
- Gilmour, M. W., Thomson, N. R., Sanders, M., Parkhill, J., and Taylor, D. E. (2004). The complete nucleotide sequence of the resistance plasmid R478: defining the backbone components of incompatibility group H conjugative plasmids through comparative genomics. *Plasmid* 52, 182–202. doi: 10.1016/j.plasmid.2004.06.006

- Gworek, B., Dmuchowski, W., Baczewska, A. H., Bragoszewska, P., Bemowska-Kalabun, O., and Wrzosek-Jakubowska, J. (2017). Air contamination by mercury, emissions and transformations—a review. *Water Air Soil Pollut.* 228:123. doi: 10.1007/s11270-017-3311-y
- Hasman, H., and Aarestrup, F. M. (2002). tcrB, a gene conferring transferable copper resistance in *Enterococcus faecium*: occurrence, transferability, and linkage to macrolide and glycopeptide resistance. *Antimicrob. Agents Chemother.* 46, 1410–1416. doi: 10.1128/aac.46.5.1410-1416.2002
- Hayashi, S., Abe, M., Kimoto, M., Furukawa, S., and Nakazawa, T. (2000). The dsbA-dsbB disulfide bond formation system of *Burkholderia cepacia* is involved in the production of protease and alkaline phosphatase, motility, metal resistance, and multi-drug resistance. *Microbiol. Immunol.* 44, 41–50. doi: 10.1111/j.1348-0421.2000.tb01244.x
- Hobman, J. L., and Crossman, L. C. (2014). Bacterial antimicrobial metal ion resistance. *J. Med. Microbiol.* 64(Pt 5), 471–497. doi: 10.1099/jmm.0.023036-0
- Huang, Z., Lu, Q., Wang, J., Chen, X., Mao, X., and He, Z. (2017). Inhibition of the bioavailability of heavy metals in sewage sludge biochar by adding two stabilizers. *PLoS One* 12:e0183617. doi: 10.1371/journal.pone.0183617
- Ianeva, O. D. (2009). [Mechanisms of bacteria resistance to heavy metals]. *Mikrobiol. Z* 71, 54–65.
- International Agency for Research on Cancer [IARC] (1990). *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Chromium, Nickel and Welding*. Lyon: IARC.
- Jagadeesan, B., Gerner-Smidt, P., Allard, M. W., Leuillet, S., Winkler, A., Xiao, Y., et al. (2019). The use of next generation sequencing for improving food safety: translation into practice. *Food Microbiol.* 79, 96–115. doi: 10.1016/j.fm.2018.11.005
- Jan, A. T., Azam, M., Siddiqui, K., Ali, A., Choi, I., and Haq, Q. M. R. (2015). Heavy metals and human health: mechanistic insight into toxicity and counter defense system of antioxidants. *Int. J. Mol. Sci.* 16, 29592–29630. doi: 10.3390/ijms161226183
- Jaroslawiecka, A., and Piotrowska-Seget, Z. (2014). Lead resistance in microorganisms. *Microbiology* 160(Pt 1), 12–25. doi: 10.1099/mic.0.070284-0
- Jarup, L. (2003). Hazards of heavy metal contamination. *Br. Med. Bull.* 68, 167–182. doi: 10.1093/bmb/ldg032
- Joutey, N. T., Sayel, H., Bahafid, W., and El Ghachtouli, N. (2015). Mechanisms of hexavalent chromium resistance and removal by microorganisms. *Rev. Environ. Contam. Toxicol.* 233, 45–69. doi: 10.1007/978-3-319-10479-9\_2
- Khromykh, A., and Solomon, B. D. (2015). The benefits of whole-genome sequencing now and in the future. *Mol. Syndromol.* 6, 108–109. doi: 10.1159/000438732
- Kim, R. Y., Yoon, J. K., Kim, T. S., Yang, J. E., Owens, G., and Kim, K. R. (2015). Bioavailability of heavy metals in soils: definitions and practical implementation—a critical review. *Environ. Geochem. Health* 37, 1041–1061. doi: 10.1007/s10653-015-9695-y
- Kourtesi, C., Ball, A. R., Huang, Y.-Y., Jachak, S. M., Vera, D. M., Khondkar, P., et al. (2013). Microbial efflux systems and inhibitors: approaches to drug discovery and the challenge of clinical implementation. *Open Microbiol. J.* 7, 34–52. doi: 10.2174/1874285801307010034
- Lee, S. M., Grass, G., Rensing, C., Barrett, S. R., Yates, C. J., Stoyanov, J. V., et al. (2002). The Pco proteins are involved in periplasmic copper handling in *Escherichia coli*. *Biochem. Biophys. Res. Commun.* 295, 616–620. doi: 10.1016/s0006-291x(02)00726-x
- Leedjävär, A., Ivask, A., and Virta, M. (2007). Interplay of different transporters in the mediation of divalent heavy metal resistance in *Pseudomonas putida* KT2440. *J. Bacteriol.* 190, 2680–2689. doi: 10.1128/jb.01494-07
- Levinson, H. S., Mahler, I., Blackwelder, P., and Hood, T. (1996). Lead resistance and sensitivity in *Staphylococcus aureus*. *FEMS Microbiol. Lett.* 145, 421–425. doi: 10.1111/j.1574-6968.1996.tb08610.x
- Li, X. Z., Plesiat, P., and Nikaido, H. (2015). The challenge of efflux-mediated antibiotic resistance in Gram-negative bacteria. *Clin. Microbiol. Rev.* 28, 337–418. doi: 10.1128/CMR.00117-14
- Lima, A. I. G., Corticeiro, S. C., and de Almeida Paula Figueira, E. M. (2005). *Glutathione-Mediated Cadmium Sequestration in Rhizobium leguminosarum*. Amsterdam: Elsevier.
- Lima de Silva, A. A., de Carvalho, M. A., de Souza, S. A., Dias, P. M., da Silva Filho, R. G., de Meirelles Saramago, C. S., et al. (2012). Heavy metal tolerance (Cr, Ag AND Hg) in bacteria isolated from sewage. *Braz. J. Microbiol.* 43, 1620–1631. doi: 10.1590/S1517-838220120004000047
- Lin, J., Michel, L. O., and Zhang, Q. (2002). CmeABC functions as a multidrug efflux system in *Campylobacter jejuni*. *Antimicrob. Agents Chemother.* 46, 2124–2131. doi: 10.1128/aac.46.7.2124-2131.2002
- Manchanda, V., Sanchaita, S., and Singh, N. (2010). Multidrug resistant acinetobacter. *J. Glob. Infect. Dis.* 2, 291–304. doi: 10.4103/0974-777X.68538
- Mancilla-Rojano, J., Castro-Jaimes, S., Ochoa, S. A., Bobadilla Del Valle, M., Luna-Pineda, V. M., Bustos, P., et al. (2019). Whole-genome sequences of five *Acinetobacter baumannii* strains from a child with leukemia M2. *Front. Microbiol.* 10:132. doi: 10.3389/fmicb.2019.00132
- Maret, W. (2015). Analyzing free zinc(II) ion concentrations in cell biology with fluorescent chelating molecules. *Metallomics* 7, 202–211. doi: 10.1039/c4mt00230j
- Markov, D., Naryshkina, T., Mustae, A., and Severinov, K. (1999). A zinc-binding site in the largest subunit of DNA-dependent RNA polymerase is involved in enzyme assembly. *Genes Dev.* 13, 2439–2448. doi: 10.1101/gad.13.18.2439
- Mata, M. T., Baquero, F., and Perez-Diaz, J. C. (2000). A multidrug efflux transporter in *Listeria monocytogenes*. *FEMS Microbiol. Lett.* 187, 185–188. doi: 10.1111/j.1574-6968.2000.tb09158.x
- Matar, G. M., Gay, E., Cooksey, R. C., Elliott, J. A., Heneine, W. M., Uwaydah, M. M., et al. (1992). Identification of an epidemic strain of *Acinetobacter baumannii* using electrophoretic typing methods. *Eur. J. Epidemiol.* 8, 9–14. doi: 10.1007/bf02427385
- Miranda, A. T., Gonzalez, M. V., Gonzalez, G., Vargas, E., Campos-Garcia, J., and Cervantes, C. (2005). Involvement of DNA helicases in chromate resistance by *Pseudomonas aeruginosa* PAO1. *Mutat. Res.* 578, 202–209. doi: 10.1016/j.mrfmmm.2005.05.018
- Mirzaei, N., Kafizadeh, F., and Kargar, M. (2008). Isolation and identification of mercury resistant bacteria from Kor river, Iran. *J. Biol. Sci.* 8, 935–939. doi: 10.3923/jbs.2008.935.939
- Morais, P. V., Branco, R., and Francisco, R. (2011). Chromium resistance strategies and toxicity: what makes *Ochrobactrum tritici* 5bv11 a strain highly resistant. *Biometals* 24, 401–410. doi: 10.1007/s10534-011-9446-1
- Naik, M. M., Pandey, A., and Dubey, S. K. (2012). *Pseudomonas aeruginosa* strain WI-1 from Mandovi estuary possesses metallothionein to alleviate lead toxicity and promotes plant growth. *Ecotoxicol. Environ. Saf.* 79, 129–133. doi: 10.1016/j.jecoen.2011.12.015
- Navy U. S. (2008). *Hawaii Range Complex: Environmental Impact Statement*. Evanston, IL: Northwestern University.
- Ndagi, U., Mhlongo, N., and Soliman, M. E. (2017). Metal complexes in cancer therapy – an update from drug design perspective. *Drug Des. Dev. Ther.* 11, 599–616. doi: 10.2147/DDDT.S119488
- Nies, D. H. (1999). Microbial heavy-metal resistance. *Appl. Microbiol. Biotechnol.* 51, 730–750. doi: 10.1007/s002530051457
- Nies, D. H. (2016). The biological chemistry of the transition metal "transportome" of *Cupriavidus metallidurans*. *Metallomics* 8, 481–507. doi: 10.1039/c5mt00320b
- Olafson, R. W., McCubbin, W. D., and Kay, C. M. (1988). Primary- and secondary-structural analysis of a unique prokaryotic metallothionein from a *Synechococcus* sp. cyanobacterium. *Biochem. J.* 251, 691–699. doi: 10.1042/bj2510691
- Osredkar, J., and Šuštar, N. (2011). Copper and zinc, biological role and significance of copper/zinc imbalance. *J. Clin. Toxicol.* S3-001, 1–19.
- Paez-Espino, D., Tamames, J., de Lorenzo, V., and Canovas, D. (2009). Microbial responses to environmental arsenic. *Biometals* 22, 117–130. doi: 10.1007/s10534-008-9195-y
- Pal, C., Asiani, K., Arya, S., Rensing, C., Stekel, D. J., Larsson, G. J. D., et al. (2017). Metal resistance and its association with antibiotic resistance. *Adv. Microb. Physiol.* 70, 261–313. doi: 10.1016/bs.ampbs.2017.02.001
- Panda, J., and Sarkar, P. (2012). Isolation and identification of chromium-resistant bacteria: test application for prevention of chromium toxicity in plant. *J. Environ. Sci. Health A Tox. Hazard. Subst. Environ. Eng.* 47, 237–244. doi: 10.1080/10934529.2012.640895
- Parks, J. M., Johs, A., Podar, M., Bridou, R., Hurt, R. A., Smith, S. D., et al. (2013). The genetic basis for bacterial mercury methylation. *Science* 339, 1332–1335. doi: 10.1126/science.1230667



- Pérez-Valdespino, A., Celestino-Mancera, M., Lorena Villegas-Rodríguez, V., and Curiel-Quesada, E. (2013). Characterization of mercury-resistant clinical *Aeromonas* species. *Braz. J. Microbiol.* 44, 1279–1283. doi: 10.1590/s1517-83822013000400036
- Perron, K., Caille, O., Rossier, C., Delden, C. V., Dumas, J.-L., and Köhler, T. (2004). CzcR-CzcS, a two-component system involved in heavy metal and carbapenem resistance in *Pseudomonas aeruginosa*. *J. Biol. Chem.* 279, 8761–8768. doi: 10.1074/jbc.m312080200
- Pradhan, S., Singh, N., Bhagwat Prasad, R., and Thatoi, H. N. (2016). Bacterial chromate reduction: a review of important genomic, proteomic, and bioinformatic analysis. *Crit. Rev. Environ. Sci. Technol.* 46, 1659–1703. doi: 10.1080/10643389.2016.1258912
- Radke, B., Jewell, L., Piketh, S., and Namieśnik, J. (2014). Arsenic-based warfare agents: production, use, and destruction. *Crit. Rev. Environ. Sci. Technol.* 44, 1525–1576. doi: 10.1080/10643389.2013.782170
- Risher, J., and Rob, D. (1999). *Toxic Substances Portal- Mercury*. Available at: <https://www.atsdr.cdc.gov/toxprofiles/tp46.pdf> (accessed March 1999).
- Rizzotto, M. (2012). *Metal Complexes as Antimicrobial Agents*. London: Intechopen.
- Rouch, D. A., Lee, B. T., and Morby, A. P. (1995). Understanding cellular responses to toxic agents: a model for mechanism-choice in bacterial metal resistance. *J. Ind. Microbiol.* 14, 132–141. doi: 10.1007/bf01569895
- Rowlatt, J. (2014). *Tungsten: The Perfect Metal for Bullets and Missiles*. London: BBC.
- Ruan, Z., and Feng, Y. (2016). BacWGSTdb, a database for genotyping and source tracking bacterial pathogens. *Nucleic Acids Res.* 44, D682–D687. doi: 10.1093/nar/gkv1004
- Savabieasfahani, M., Ali, S. S., Bacho, R., Savabi, O., and Alsabbak, M. (2016). Prenatal metal exposure in the Middle East: imprint of war in deciduous teeth of children. *Environ. Monit. Assess.* 188:505. doi: 10.1007/s10661-016-5491-0
- Scott, P., Deye, G., Srinivasan, A., Murray, C., Moran, K., Hulten, E., et al. (2007). An outbreak of multidrug-resistant *Acinetobacter baumannii-calcoaceticus* complex infection in the us military health care system associated with military operations in Iraq. *Clin. Infect. Dis.* 44, 1577–1584. doi: 10.1086/518170
- Seiler, C., and Berendonk, T. (2012). Heavy metal driven co-selection of antibiotic resistance in soil and water bodies impacted by agriculture and aquaculture. *Front. Microbiol.* 3:399. doi: 10.3389/fmicb.2012.00399
- Sharma, J., Shamim, K., Dubey, S. K., and Meena, R. M. (2017). Metallothionein assisted periplasmic lead sequestration as lead sulfite by *Providencia vermicola* strain SJ2A. *Sci. Total Environ.* 579, 359–365. doi: 10.1016/j.scitotenv.2016.11.089
- Shin, M. N., Shim, J., You, Y., Myung, H., Bang, K. S., Cho, M., et al. (2012). Characterization of lead resistant endophytic *Bacillus* sp. MN3-4 and its potential for promoting lead accumulation in metal hyperaccumulator *Alnus firma*. *J. Hazard. Mater.* 199–200, 314–320. doi: 10.1016/j.jhazmat.2011.11.010
- Silveira, E., Freitas, A. R., Antunes, P., Barros, M., Campos, J., Coque, T. M., et al. (2014). Co-transfer of resistance to high concentrations of copper and first-line antibiotics among *Enterococcus* from different origins (humans, animals, the environment and foods) and clonal lineages. *J. Antimicrob. Chemother.* 69, 899–906. doi: 10.1093/jac/dkt479
- Silver, S., and Phung le, T. (2005). A bacterial view of the periodic table: genes and proteins for toxic inorganic ions. *J. Ind. Microbiol. Biotechnol.* 32, 587–605. doi: 10.1007/s10295-005-0019-6
- Singh, R., Gautam, N., Mishra, A., and Gupta, R. (2011). Heavy metals and living systems: an overview. *Indian J. Pharmacol.* 43, 246–253. doi: 10.4103/0253-7613.81505
- Singh, S. K., Roberts, S. A., McDevitt, S. F., Weichsel, A., Wildner, G. F., Grass, G. B., et al. (2011). Crystal structures of multicopper oxidase CueO bound to copper(I) and silver(I): functional role of a methionine-rich sequence. *J. Biol. Chem.* 286, 37849–37857. doi: 10.1074/jbc.M111.293589
- Smeaton, C. M., Fryer, B. J., and Weisener, C. G. (2009). Intracellular precipitation of Pb by *Shewanella putrefaciens* CN32 during the reductive dissolution of Pb-jarosite. *Environ. Sci. Technol.* 43, 8086–8091. doi: 10.1021/es901629c
- Subramanian, S. (2018). Studies on the biosorption of Pb(II) ions using *Pseudomonas putida*. *Sep. Sci. Technol.* 53, 2550–2562. doi: 10.1080/01496395.2018.1464025
- Szivač, I., Behra, R., and Sigg, L. (2009). Metal-induced reactive oxygen species production in *Chlamydomonas reinhardtii* (chlorophyceae). *J. Phycol.* 45, 427–435. doi: 10.1111/j.1529-8817.2009.00663.x
- Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., and Sutton, D. J. (2012). Heavy metal toxicity and the environment. *Exp. Suppl.* 101, 133–164.
- Tong, M. J. (1972). Septic complications of war wounds. *JAMA* 219, 1044–1047. doi: 10.1001/jama.219.8.1044
- Trevors, J. T., and Oddie, K. M. (1986). R-plasmid transfer in soil and water. *Can. J. Microbiol.* 32, 610–613. doi: 10.1139/m86-114
- Williams, C. L., Neu, H. M., Gilbreath, J. J., Michel, S. L., Zurawski, D. V., and Merrell, D. S. (2016). Copper resistance of the emerging pathogen *Acinetobacter baumannii*. *Appl. Environ. Microbiol.* 82, 6174–6188. doi: 10.1128/aem.01813-16
- Wilson, A., Fox, E. M., Fegan, N., and Kurtboke, D. I. (2019). Comparative genomics and phenotypic investigations into antibiotic, heavy metal, and disinfectant susceptibilities of *Salmonella enterica* strains isolated in Australia. *Front. Microbiol.* 10:1620. doi: 10.3389/fmicb.2019.01620
- Wireman, J., Liebert, C. A., Smith, T., and Summers, A. O. (1997). Association of mercury resistance with antibiotic resistance in the gram-negative fecal bacteria of primates. *Appl. Environ. Microbiol.* 63, 4494–4503. doi: 10.1128/aem.63.11.4494-4503.1997
- Wong, P. T. S., Chau, Y. K., and Luxon, P. L. (1975). Methylation of lead in the environment. *Nature* 253, 263–264. doi: 10.1038/253263a0
- Xia, X., Li, J., Zhou, Z., Wang, D., Huang, J., and Wang, G. (2018). High-quality-draft genome sequence of the multiple heavy metal resistant bacterium *Pseudaminobacter manganicus* JH-7(T). *Stand. Genomic Sci.* 13:29. doi: 10.1186/s40793-018-0330-2
- Yang, J., Rawat, S., Stemmler, T. L., and Rosen, B. P. (2010). Arsenic binding and transfer by the ArsD As(III) metallochaperone. *Biochemistry* 49, 3658–3666. doi: 10.1021/bi100026a
- Yu, Z., Gunn, L., Wall, P., and Fanning, S. (2017). Antimicrobial resistance and its association with tolerance to heavy metals in agriculture production. *Food Microbiol.* 64, 23–32. doi: 10.1016/j.fm.2016.12.009

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Bazzi, Abou Fayad, Nasser, Haraoui, Dewachi, Abou-Sitta, Nguyen, Abara, Karah, Landecker, Knapp, McEvoy, Zaman, Higgins and Matar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.